arXiv:hep-ph/0010058v1 6 Oct 2000

Flavour–dependent SUSY phases and CP asymmetry in $B \to X_s\gamma$ decays

D. Bailin and S. Khalil
Centre for Theoretical Physics, University of Sussex, Brighton BN1 9QJ, U. K.

(March 2, 2009)

We study the direct CP asymmetry in $B \to X_s\gamma$ decay in SUSY models with non–universal A–terms. We show that the flavour–dependent phases of the A–terms, unlike the flavour–independent ones, can give rise too large contribution to the CP asymmetry while respecting the experimental bounds on the branching ratio of $B \to X_s\gamma$ decay and the electric dipole moments of the electron and neutron. The CP asymmetry of this decay is predicted to be much larger than the standard model prediction in a wide region of the parameter space. In particular, it can be of order $10-15\%$ which can be accessible at B factories.

PACS numbers:12.60.Jv, 11.30.Er, 13.20.He SUSX-TH-00-015

Direct CP asymmetry in the inclusive radiative decay $B \to X_s\gamma$ is measured by the quantity

$$A_{CP}^{b\to s\gamma} = \frac{\Gamma(\bar{B} \to X_s\gamma) - \Gamma(B \to X_{\bar{s}}\gamma)}{\Gamma(\bar{B} \to X_s\gamma) + \Gamma(B \to X_{\bar{s}}\gamma)}. \quad (1)$$

The Standard Model (SM) prediction for this asymmetry is very small, less than 1%. Thus, the observation of sizeable asymmetry in the decay $B \to X_s\gamma$ would be a clean signal of new physics.

The most recent result reported by CLEO collaboration for the CP asummetry in these decays is [1]

$$-9\% < A_{CP}^{b\to s\gamma} < 42\%\ , \quad (2)$$

and it is expected that the measurements of $A_{CP}^{b\to s\gamma}$ will be improved in the next few years at the B–factories.

Supersymmetric predictions for $A_{CP}^{b\to s\gamma}$ are strongly dependent on the flavour structure of the soft breaking terms. It was shown that in the universal case, as in the minimal supergravity models, the prediction of the asymmetry is less than 2%, since in this case the electric dipole moments (EDM) of the electron and neutron constrain the SUSY CP–violating phases to be very small [2,3]. Furthermore, it is also known that in this case, one can not get any sizeable SUSY contribution to the CP–violating observables, ε and ε'/ε [4–7]. The impact of flavour structure in the non–degenerate A–terms, which is expected in string inspired models, has been studied [6–8]. In these works, it was emphasized that the flavour–dependent SUSY CP–violating phases are not essentially constrained by the EDMs and can lead to large contributions to the observed CP–violating phenomena in the kaon system and, in particular, to the direct CP violation measured by ε'/ε.

In this letter, we explore the effect of these large flavour–dependent phases on inducing a direct CP violation in $B \to X_s\gamma$ decay. We will show that the values of the asymmetry $A_{CP}^{b\to s\gamma}$ in this class of models are much larger than the SM prediction in a wide region of the parameter space allowed by experiments, namely the EDM experimental limits and the bounds on the branching ratio of $B \to X_s\gamma$. The enhancement of $A_{CP}^{b\to s\gamma}$ is due to the important contributions from gluino–mediated diagrams, in this scenario, in addition to the usual chargino and charged Higgs contributions.

The effective Hamiltonian for $b \to s\gamma$ decay is given by

$$H_{eff} = -\frac{4G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_{i=1}^{8} C_i(\mu_b) Q_i(\mu_b) \quad (3)$$

where $Q_i(\mu_b)$ and $C_i(\mu_b)$ represent an operator for the $\Delta B = 1$ transition and its Wilson coefficient, respectively, evaluated at the renormalization scale $\mu_b \simeq \mathcal{O}(m_b)$. The relevant operators for $b \to s\gamma$ decay are given by

$$\begin{aligned} Q_2 &= \bar{s}_L \gamma^\mu c_L\ \bar{c}_L \gamma_\mu b_L, \\ Q_7 &= \frac{e}{16\pi^2} m_b \bar{s}_L \sigma^{\mu\nu} b_R F_{\mu\nu}, \\ Q_8 &= \frac{g_s}{16\pi^2} m_b \bar{s}_L \sigma^{\mu\nu} T^a b_R G^a_{\mu\nu}. \end{aligned} \quad (4)$$

The expression for the asymmetry $A_{CP}^{b\to s\gamma}$, corrected to next–to–leading order is given by [9]

$$\begin{aligned} A_{CP}^{b\to s\gamma} = \frac{4\alpha_s(m_b)}{9|C_7|^2} \Big\{ & \Big[\frac{10}{9} - 2z\ (v(z) + b(z,\delta))\Big] Im[C_2 C_7^*] \\ & + Im[C_7 C_8^*] + \frac{2}{3} z\ b(z,\delta) Im[C_2 C_8^8] \Big\}, \end{aligned} \quad (5)$$

where $z = m_c^2/m_b^2$. The functions $v(z)$ and $b(z,\delta)$ can be found in Ref. [9]. The parameter δ is related to the experimental cut on the photon energy, $E_\gamma > (1-\delta) m_b/2$, which is assumed to be 0.9. We neglect the very small effect of the CP–violating phase in the CKM matrix.

The SUSY contributions to the Wilson coefficients $C_{7,8}$ are obtained by calculating the $b \to s\gamma$ and $b \to sg$ amplitudes at the electroweak (EW) scale respectively. The leading–order contributions to these amplitudes are given by the 1–loop magnetic-dipole and chromomagnetic dipole penguin diagrams respectively, mediated by charged Higgs boson, chargino, gluino, and neutralino exchanges. As pointed out in Ref. [10], SUSY models with non–universal A–terms may induce non–negligible contributions to the dipole operators $\tilde{Q}_{7,8}$ which have opposite chirality to $Q_{7,8}$. In the MSSM these contributions are suppressed by terms of order $\mathcal{O}(m_s/m_b)$ due to the universality of the A–terms. However, in our case we should take them into account. Denoting by $\tilde{C}_{7,8}$ the Wilson coefficients multiplying the new operators $\tilde{Q}_{7,8}$ the expression for the asymmetry in Eq.(5) will be modified by making the replacement

$$C_i C_j^* \to C_i C_j^* + \tilde{C}_i \tilde{C}_j^*. \tag{6}$$

The expressions for $\tilde{C}_{7,8}$ are given in Ref. [10] and $\tilde{C}_2 = 0$ (there is no operator similar to Q_2 containing right–handed quark fields).

In this case the general parametrization of the branching ratio in terms of the new physics contributions takes the form [10,11]

$$\begin{aligned} BR(B \to X_s\gamma) = (3.29 \pm 0.33) \times 10^{-4} \Bigg(& 1 + 0.622 Re[\xi_7] \\ & + 0.090\big(\, |\xi_7|^2 + |\tilde{\xi}_7|^2 \, \big) + 0.066 \; Re[\xi_8] + 0.019\big(\; Re[\xi_7 \xi_8^*] \\ & + Re[\tilde{\xi}_7 \tilde{\xi}_8^*] \, \big) + 0.002\big(\, |\xi_8|^2 \; + \; |\tilde{\xi}_8|^2 \, \big) \Bigg) , \end{aligned} \tag{7}$$

where $\xi_{7,8} = \big(C_{7,8} - C_{7,8}^{SM}\big)/C_{7,8}^{SM}$ and $\tilde{\xi}_{7,8} = \tilde{C}_{7,8}/C_{7,8}^{SM}$. $C_{7,8}$ include the total contribution while $C_{7,8}^{SM}$ contains only the SM ones.

Note that including these modifications (6) may enhance the branching ratio of $B \to X_s\gamma$ and reduce the CP asymmetry, since $|C_7|^2$ is replaced by $|C_7|^2 + |\tilde{C}_7|^2$ in the denominator of Eq.(5). If so, neglecting this contribution could lead to an incorrect conclusion.

As already mentioned, usually the SUSY trilinear terms are defined at the high energy scale to be proportional to their associated Yukawa matrix, *i.e.* $Y_{ij}^A = A_0 Y_{ij}$ and A_0 is the same for all Yukawa couplings. Despite the simplicity of this assumption, it is a very peculiar case and there exist classes of model inspired by string theory that lead to non–universal A–terms. Moreover, this non–universality is crucial in order to generate the experimentally observed CP violation ε and ε'/ε even with the phase δ_{CKM} vanishing [6–8].

Motivated by these models [7] we parameterize the trilinear matrices Y_d^A as $(Y_d^A)_{ij} = A_{ij}^d Y_{ij}^d$, where A_{ij}^d is given by

$$A_{ij}^d = \begin{pmatrix} a & a & b \\ a & a & b \\ b & b & c \end{pmatrix}. \tag{8}$$

The entries a, b and c are complex and of order the gravitino mass, $\mathcal{O}(m_{3/2})$. In general A^d and A^u have different structure. Here we assume for simplicity that $A^d = A^u$. The results, as we will show, are sensitive only to the non–universality between the entries A_{ij}. Also we assume that the scalar masses and the gaugino masses are universal and of order $m_{3/2}$ too. So that after rotating the phase of the gaugino masses we have three phases in the A–sector: ϕ_a, ϕ_b and ϕ_c, which are the relative phases between the gaugino phase and the original phases of the A_{ij} entries. It is important to note that the EDM limits on the electron and the neutron essentially constrain the phase of the entry A_{11} (ϕ_a) and the other two phases are much less constrained. These two phases gives large SUSY contribution to the CP observables.

We found that the flavour–independent phase ϕ_c gives a large contribution to $\tilde{C}_7$ and $\tilde{C}_8$. This can simply understood by using the mass insertion, the gluino contributions of $\tilde{C}_7$ and $\tilde{C}_8$ are proportional to $(\delta_{LR}^d)_{23} \simeq (S_{D_L} Y_d^{A*} S_{D_R}^\dagger)_{23}/m_{\tilde{q}}^2$ and $(\delta_{RL}^d)_{23} \simeq (S_{D_R} Y_d^A S_{D_L}^\dagger)_{23}/m_{\tilde{q}}^2$, where $S_{D_R} Y_d S_{D_L}^\dagger = Y_d^{diag}$. It is obvious that the entries A_{33} could give significant contribution to $(\delta_{LR}^d)_{23}$ and $(\delta_{RL}^d)_{23}$. As explained above, enhancing the Wilson coefficients $\tilde{C}_{7,8}$ increases the value of the branching ratio and decreases the value of the CP asymmetry.

Indeed, we found that for small values ϕ_c the branching ratio is very much enhanced and becomes above the experimental upper bound, and the CP asymmetry is reduced a lot. Thus ϕ_c is constrained to be very small ($\lesssim \mathcal{O}(10^{-1})$), and since we are interested in studying scenarios in which the CP asymmetry is sizeable, we assume that ϕ_c is zero*. In other words, the flavour–diagonal phases have to be aligned with the gaugino phase. In fact, this would be true in some string inspired models where the dilaton vacuum expectation value (VEV) gives the main contribution to the gaugino masses and the diagonal entries of the A–terms, while the VEVs of the moduli fields give the contribution to the off–diagonal elements of A–terms [7].

The effect of the other flavour–independent phase, ϕ_a on the $A_{CP}^{b\to s\gamma}$ and $BR(b \to s\gamma)$ is found to be very small. So any constraint on this phase from the EDM is not a problem in this class of models.

It is worth mentioning that the textures of the A–terms that derived from type I string inspired models are not favoured here since they can not enhance the CP asymmetry over that in the minimal supergravity models. These textures have the following form [8]

$$A_{ij} = \begin{pmatrix} a & b & c \\ a & b & c \\ a & b & c \end{pmatrix}, \tag{9}$$

which means that we can write the parameters Y_{ij}^A in matrix notation as

---

*In any case, we expect that the EDM of the mecury atom provides a severe constraint on this phase [12]

$$Y^A = Y \ . \ \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} .$$

Thus, working in the SCKM basis where the unitary matrices $S_{L,R}$ are obtained by superfield rotation so that the quark mass matrices are diagonal, the trilinear terms become $Y_{diag}.A'$ where A' is given by $A' = S_L A S_L^\dagger$. This result is very similar to the case of the universal A–terms. Since the Yukawa couplings preserve the diagonality once this basis is chosen at the GUT scale, the effect of the off–diagonal element in this model is diluted. Moreover, as seen from Eq.(9), the phases of the elements A_{22} and A_{33} (which are constrained by the EDM and the $BR(b \to s\gamma)$) are the same phases of the off–diagonal elements A_{32} and A_{23} respectively. Therefore, this kind of texture does not give a significant contribution to the CP asymmetry in the $B \to X_s\gamma$, as pointed out in Ref. [13], although it gives important contribution to the CP–violating effects in the kaon system, as shown in Ref. [8].

In the class of models we consider here, Eq.(8), the relevant and important phase for the CP asymmetry is the phase of the off–diagonal elements ϕ_b. In Fig.1 we show the dependence of $A_{CP}^{b\to s\gamma}$ on ϕ_b for $m_{3/2} = 150$ GeV and $\tan\beta = 3$ and 10. The non–universality among $|a|$, $|b|$ and $|c|$ is crucial to get these results. Large $|c|$ is required for obtaining the $BR(b \to s\gamma)$ values within the experimental limits, especially at low $\tan\beta$. This is because large $|c| \equiv A_t$ leads to large off–diagonal elements in the stop mass matrix. Hence the splitting of the two stop mass eigenstates becomes large and this enhances the chargino contribution which decreases the total branching ratio [14].

Therefore, for $\tan\beta = 10$ we fixed $|c| \simeq \sqrt{3} m_{3/2}$ and for $\tan\beta = 3$ we used $|c| \simeq 3\ m_{3/2}$ otherwise the values of the branching ratio of $B \to X_s\gamma$ decay are larger than 4×10^{-4}. Also we observe that $|b| < |c|$ is favoured by large CP asymmetry $A_{CP}^{b\to s\gamma}$.

We see from Fig.1 that the CP asymmetry $A_{CP}^{b\to s\gamma}$ can be as large as ±15%, which can be accessible at the B–factories. Also this result does not require a light chargino as in the case considered in Ref. [15].

It is important to emphasize that the gluino contribution in this model gives the dominant contribution to the CP asymmetry $A_{CP}^{b\to s\gamma}$. We found that although the real parts of the gluino contributions to both of $C_{7,8}$ and $\tilde{C}_{7,8}$ are smaller than the real parts of the other contributions (but not negligible as in the case of universal A–terms), their imaginary parts are dominant and give with the imaginary parts of the chargino contribution the main contributions to $A_{CP}^{b\to s\gamma}$. It is clear that these contributions vanish for ϕ_b equal to a multiple of π and $A_{CP}^{b\to s\gamma}$ in this case is identically zero as Fig.1 shows.



FIG. 1. CP asymmetry $A_{CP}^{b\to s\gamma}$ as a function of the flavour–dependent phase ϕ_b, for $m_{3/2} \simeq 150$ GeV and $\tan\beta = 3$ and 10.

In Fig.2 we show the correlation between the CP asymmetry $A_{CP}^{b\to s\gamma}$ and the branching ratio $BR(B \to X_s\gamma)$. We scan over the following region of the parameter space:

$$\tan\beta \simeq 2{-}40, \quad m_0,\ m_{1/2},\ |a|,\ |b|,\ |c| \simeq 100{-}500\,GeV,$$

where m_0 is the common scalar mass and $m_{1/2}$ is the common gaugino masses. Assuming non–universal scalar and gaugino masses could also help in enhancing the CP asymmetry, however, here we assume these masses are universal to maximizing the effect of the non–universality of the trilinear couplings.

The phases ϕ_a and ϕ_c are assumed to be zero while ϕ_b is fixed to be $\pi/2$ which gives, as shown in Fig.1, nearly maximum CP asymmetry. The low energy SUSY spectrum is calculated by running these soft SUSY breaking terms from the GUT scale ($\simeq 2 \times 10^{16}$ GeV) to the EW scale ($\simeq M_Z$) and imposing all the usual constraints, like the electroweak symmetry breaking, absence of colour and electric charge breaking minima, experimental bound on the SUSY particles, etc.

From this figure we see that there is a wide region of the parameter space where the decay rate asymmetry of $B \to X_s\gamma$ can be large. We also note that large values of CP asymmetry $A_{CP}^{b\to s\gamma}$ prefer small values for the branching ratio $BR(B \to X_s\gamma)$. This correlation is also found in Ref. [3]. We fixed the sign of μ to be positive (in our convention, where the sign of μ in chargino mass matrix is positive and in neutralino mass matrix is negative) since $\mu < 0$ leads to large values of $BR(B \to X_s\gamma)$ and almost the whole range of parameter space is excluded.

In summary, we have considered the possible supersymmetric contribution to CP asymmetry in the inclusive $B \to X_s\gamma$ decay in model with non–universal A–terms. Contrary to the universal scenario, we find that the CP asymmetry in this class of models is predicted to be large



FIG. 2. Correlations of CP asymmetry $A_{CP}^{b\to s\gamma}$ with the branching ratio $BR(B \to X_s\gamma)$.

in sizeable regions of the parameter space allowed by the experimental bounds, and may be possibly to be detected at B factories We have shown that the flavour–dependent phases are crucial for this enhancing with respecting the severe bounds on the electric dipole moment of the neutron and electron.

ACKNOWLEDGMENTS

S.K. would like to thank S. Abel, F. Borzomati and M. Gomez for useful discussions. This work was supported by the PPARC.

[1] CLEO Collaboration, S. Ahmed et al., hep-ex/9908022.
[2] T. Goto, Y. Keum, T. Nihei, Y. Okada and Y. Shimizu, Phys. Lett. B460 (1999) 333;
[3] M. Aoki, G. Cho and N. Oshimo, Nucl. Phys. B554 (1999) 50.
[4] D. Demir, A. Masiero, and O. Vives, Phys. Lett. B 479 (2000) 230.
[5] S. Barr and S. Khalil, Phys. Rev. D61 (192000) 035005.
[6] S. A. Abel and J.-M. Frére, Phys. Rev. D55 (1997) 1623.
[7] S. Khalil, T. Kobayashi and A. Masiero, Phys. Rev. D60 (1999) 075003; S. Khalil and T. Kobayashi, Phys. Lett. B460 (1999) 341.
[8] S. Khalil, T. Kobayashi and O. Vives , Nucl. Phys. B580 (2000) 275.
[9] A. Kagan and M. Neuber, Phys. Rev. D58 (1998) 094012.
[10] E. Gabrielli, S. Khalil, E. Torrente-Lujan, hep-ph/0005303 , to appear in Nucl. Phys. B.
[11] A. Kagan and M. Neubert, Eur. Phys. J. C 7 (1999) 5.
[12] S. Abel, D. Bailin and S. Khalil, “Complete Analysis of EDM: Cancellation Mechanism vs. Non–universality” (in preparation).
[13] A. Masiero and O. Vives , hep-ph/0007320.
[14] S. Khalil , A. Masiero and Q. Shafi , Phys. Rev. D56 (1997) 5754 and references therein.
[15] S. Baek and P. Ko, Phys. Rev. Lett. 83 (1999) 488.